

## IMPACTS OF CLIMATE CHANGE ON MARINE MAMMALS

Peter G.H. Evans<sup>1</sup>, Ian L. Boyd<sup>2</sup> & Colin D. MacLeod<sup>3</sup>

### Organisation(s) represented (To include full contact details)

<sup>1</sup> **Sea Watch Foundation (SWF)**, Cynifryn, Abershore, Llanfaglan, Caernarfon, Gwynedd LL54 5RA. Tel: 01286-672071/0776-556-6102.

Email: [peter.evans@seawatchfoundation.org.uk](mailto:peter.evans@seawatchfoundation.org.uk);

Web: [www.seawatchfoundation.org.uk](http://www.seawatchfoundation.org.uk)

<sup>2</sup> **Sea Mammal Research Unit (SMRU)**, Gatty Marine Laboratory, University of St Andrews, St Andrews, Fife KY16 8LB. Email: [ilb@st-andrews.ac.uk](mailto:ilb@st-andrews.ac.uk);

Web: <http://smub.st-and.ac.uk>

<sup>3</sup> **University of Aberdeen**, School of Biological Sciences (Zoology), Tillydrone Avenue, Aberdeen AB24 3JG. Email: [c.d.macleod@abdn.ac.uk](mailto:c.d.macleod@abdn.ac.uk)

**Please can you provide a review (2-3 pages) which broadly synthesises current scientific understanding and debate on the impacts of climate change on **Marine Mammals** important to the UK marine environment<sup>o</sup>**

### Overview

As warm-blooded vertebrates with both physiological and behavioural mechanisms developed to respond readily to changes in their external environment, marine mammals might be expected to cope well with most environmental variance predicted from climate change. On the other hand, we would anticipate changes in the availability of their habitat (including food resources) to lead to functional responses in particular cases. The most obvious example in this context is the reduction in ice cover affecting ice-breeding polar seals such as the ringed seal, and its consequent effect upon predators like the polar bear (Stirling *et al.*, 1999; Derocher *et al.*, 2004; Ferguson *et al.*, 2005). Other physical changes to the habitat include increased sea levels modifying shallow seas. These could affect breeding bays & lagoons of coastal species like the vaquita and grey whale (IWC, 1997), and seal caves and other haul-out sites (e.g. the endangered Mediterranean monk seal – Harwood, 2001). Potentially also vulnerable are dugongs and manatees, humpback dolphins, franciscana, tucuxi, and finless porpoise all of whose diets appear to be dependent upon shallow-water species (sea grasses in the case of sirenians – see Short & Neckles, 1999; and fishes and invertebrates in the case of cetaceans (Würsig *et al.*, 2002). However, in the UK, those conditions are not likely to be important at least in the foreseeable future, and even elsewhere, it may be that new habitats become available or that species adapt within the predicted time scale.

Most likely impacts of climate change on marine mammals will be in the form of range changes through effects upon ectothermic prey. Although most marine mammals appear to have catholic diets, nevertheless in many cases species distributions fall between particular sea temperature boundaries, and this reflects the range preferences of the major prey organisms upon which they are known to feed (Bjørge, 2002).

### **What is happening now**

Recent warming of the seas around the British Isles has coincided with a northward shift in the distribution of zooplankton (Beaugrand *et al.*, 2002; Reid *et al.*, 2003) and fish species (Beare *et al.*, 2005; Perry *et al.*, 2005). Those changes are most evident near the northern or southern boundaries of the species range (Rose, 2005). The same appears to be the case for certain cetacean species, with the typical warmer water dolphins – short-beaked common dolphin and striped dolphin, recently extending their shelf sea range further north off western Britain and around into the northern North Sea (Evans *et al.*, 2003; MacLeod *et al.*, 2005). Other warm water species recorded for the first time in the UK in recent years include Blainville's beaked whale (1993) and Fraser's dolphin (1996), whilst nine out of ten strandings of pygmy sperm whale in Britain & Ireland have occurred since 1980 (Evans *et al.*, 2003). However, it is unwise to draw too many conclusions from vagrant records such as these; additional warm water species that were recorded in the UK more than fifty years ago include false killer whale (last stranding in 1935) and rough-toothed dolphin (one stranding only, in 1949).

The ability to detect long-term trends in cetaceans around the UK is limited by the paucity of effort-based sightings data before 1980. The recent multiple presence of short-beaked common dolphin schools off northern Scotland and in the northern North Sea is not entirely new, for example. There have been both strandings and sightings in that region during the 1980s (Sheldrick, 1989; Baines *et al.*, 2006), and a marked peak in North Sea strandings was reported during the 1930s (Fraser, 1946). Nevertheless, these may reflect decadal climatic cycles such as caused by the North Atlantic Oscillation (which was in a positive phase during those periods, characterised by stronger westerly winds, higher sea temperatures, and milder winters with wetter and stormier conditions (Hulme *et al.*, 2002).

Another species reported to have recently exhibited latitudinal range shifts is the white-beaked dolphin, a species of largely cold temperate to arctic waters (Evans *et al.*, 2003, MacLeod *et al.*, 2005, 2007; Baines *et al.*, 2006). On the other hand, these may not necessarily be directly related to sea temperature changes but simply reflect changes in the status of particular fish stocks that are favoured prey of the species. Stranding records have shown a significant increase in the southern North Sea since the 1960s, and the species now regularly occurs in the Southern Bight (Bakker & Smeenk 1987, Kinze *et al.* 1997; Camphuysen & Peet, 2006). Likewise, Risso's dolphins (which feed principally upon small cephalopods such as cuttlefish and octopus) are now recorded regularly in the North Sea off East Scotland and Northeast England, after being very scarce here between the 1980s and 2000 (Evans *et al.*, 2003; Sea Watch Foundation database). This emphasises the difficulties in interpreting regional changes in status, not only because there are many potential confounding effects (not least being human over-exploitation of fish or squid stocks) but also because those marine mammal species may simply be responding to regional variability in resource availability independent of climate change.

Amongst marine mammals, information on population trends, breeding success and feeding ecology is best for seals. However, at present, there is no evidence that climate change has affected either of the two UK breeding species (grey and harbour seal), although elsewhere in the world, several authors have attempted to link changes in seal population dynamics and life history parameters to climate change (Sun *et al.*, 2004, McMahon and Burton, 2005; Forcada *et al.*, 2005).

### **What could happen in the future**

Responses both at the individual and population level of marine mammal species to climate change are currently poorly understood. Making predictions about future impacts becomes even more speculative. In the last ten years, a number of marine mammal scientists have attempted to do this (Tynan & DeMaster, 1997; IWC, 1997; Würsig *et al.*, 2002; Learmonth *et al.*, 2006; Simmonds & Isaac, 2007). The main potential impacts are summarised under the following headings:

*Range shifts:* As a result of increased sea temperatures, it is thought that species will shift their ranges latitudinally. This may not necessarily result in a negative response (unless there are effects upon competitive interactions), although species with restricted distributions such as the vaquita, river dolphins, bowhead whale, narwhal, and polar seals, may be unable to change their geographic range (Tynan & DeMaster, 1997; IWC, 1997). In the UK, one might expect species like the short-beaked common and striped dolphin to occur more regularly in northern Britain, displacing the white-beaked and Atlantic white-sided dolphin. Cuvier's beaked whale may become more regular in offshore canyons, and there could be more records of warm-water vagrants to North-west Europe (e.g. Bryde's whale, pygmy sperm whale, rough-toothed dolphin, and Atlantic spotted dolphin). Baleen whales (e.g. humpback whale) that move southwards to warmer waters to winter and breed may increasingly do so within UK waters.

*Changes to physical habitat:* The melting of sea ice clearly poses a threat to those marine mammals (such as seals) that use it for hauling out or breeding, as well as to their predators (Harwood, 2001; Derocher *et al.*, 2004; Ferguson *et al.*, 2005). Changes to open water refugia in the ice may affect species like beluga and narwhal (Heide-Jørgensen & Laidre, 2004; Laidre & Heide-Jørgensen, 2005). Rising sea levels may affect shallow water species such as tucuxi, humpback dolphin, and finless porpoise, as well as those species such as the grey whale, calving in shallow coastal bays (IWC, 1997; Würsig *et al.*, 2002). In the UK, however, it is unlikely that changes to the physical habitat will affect cetaceans, although some seal haul-out / breeding locations in caves or on low lying coasts (e.g. Monach Isles for grey seals, parts of The Wash for harbour seals) may be lost or modified. Increases in storm frequency and associated wave surges could exacerbate effects. On the other hand, seals may adapt to these changes and new habitats may be created.

*Changes to the food web:* Effects of changes to community structure are probably the most difficult to predict. Changes in ocean currents and the positions of associated fronts as well as in ocean mixing, deep water production and coastal upwellings could have profound effects on biological productivity (Walther *et al.*, 2002; Stenseth *et al.*, 2002, 2004), which in turn is likely to affect top predators such as marine mammals (IWC, 1997; Bjørge, 2002; Würsig *et al.*, 2002). Species with restricted diets (such as right whales which feed largely upon copepod aggregations) are thought to be particularly vulnerable (Greene & Pershing, 2004; Leaper *et al.*, 2005). However, in most cases it is hard to determine how those changes would impact particular marine mammal species (Trites *et al.*, 2006).

More complicated scenarios include mismatches in synchrony between predator and prey, either in time or location (Edwards & Richardson, 2004). This uncoupling of phenological relationships at different trophic levels is thought to have been responsible for recent failures in sandeel recruitment in the North Sea (Arnott & Ruxton, 2002) and consequent breeding failures amongst several UK seabird species largely dependent on sandeels (Rindorf *et al.*, 2000; Frederiksen *et al.*, 2004, 2005; Wanless *et al.*, 2005; Mavor *et al.*, 2006). There has been much speculation that the recent shift in abundance of harbour porpoises from the northern to southern North Sea may be due to a shortage of sandeels, a known prey item, and that this has led to suggestions of food starvation amongst stranded porpoises (MacLeod *et al.*, 2007). On the other hand, if these observations do indeed reflect a real change, it may also be due to other prey-related changes such as the recovery of some herring stocks in the southern North Sea. At present, we have little idea to what extent species like the harbour porpoise have particular dietary preferences although high-energy shoaling fish such as herring, sprat and sandeel often form important components of porpoise diet (Santos *et al.*, 2004; Evans *et al.*, 2007).

A number of findings indicating potential effects on other marine taxa could also impact upon marine mammals through the food chain. Examples include reductions in salinity, increases in CO<sub>2</sub> and consequent decreases in pH particularly affecting cephalopods (Boyle, 1983; Pörtner *et al.*, 2004; Royal Society, 2005). Several marine mammal species feed either exclusively or to a large extent upon cephalopods.

*Susceptibility to disease and contaminants:* Global warming has been implicated in the worldwide increase in reports of diseases affecting marine organisms, including marine mammals (Harvell *et al.*, 2002; Geraci & Lounsbury, 2002; Lafferty *et al.*, 2004). Climate change has the potential to increase pathogen development and survival rates, disease transmission, and host susceptibility (Harvell *et al.*, 2002), whilst higher temperatures may stress organisms, increasing their susceptibility to some diseases (Lafferty *et al.*, 2004).

The frequency and severity of toxic algal blooms are also predicted to increase as a result of nutrient enrichment (increased rainfall and freshwater runoff) and increased temperature (Peperzak, 2003). Mass die-offs due to fatal poisonings have been reported in several marine mammal species, for example Mediterranean monk seals, California sea lions, bottlenose dolphins, and Florida manatees (Hernández *et al.*, 1998; Geraci *et al.*, 1999; Scholin *et al.*, 2000; Geraci & Lounsbury, 2002; Domingo *et al.*, 2002).

Finally, there are some suggestions that habitat degradation may occur through effects of climate upon pollutant burdens (MacDonald *et al.*, 2005).

As yet, most of the above predictions remain speculative and unsubstantiated by unequivocal evidence. This reflects the difficulties in studying marine mammals, using experimental approaches to test hypotheses, and disentangling various potential confounding variables.

**Please can you provide an executive summary (c. 200 words) that highlights the key messages from your review document**

- The impact of climate change on marine mammals remains poorly understood. As a result, there has been a great deal of speculation but without very much substantive evidence. That evidence for marine mammals is very difficult to obtain, particularly when there are often synergistic effects from anthropogenic activities.
- Most obvious impacts are loss of available habitat such as ice cover to ice-breeding pinnipeds. This already is thought to affect ringed seals and their main predator, the polar bear, in arctic regions.
- In UK waters, environmental changes will likely be reflected mainly in responses to prey distribution changes as a result of warmer sea temperatures.
- Range shifts can be observed in a number of odontocete cetacean species, but at present it is not possible to differentiate between short-term responses to regional resource variability and longer-term ones driven by climate change.
- A wide range of potential impacts of climate change can be listed, mainly operating through marine mammal prey (zooplankton, fish and cephalopod abundance and distribution changes). The relative importance of these and their likelihood of occurrence, remain unknown.

**Say what qualitative level of confidence you can place in the science for 'what is already happening' and 'what could happen in the future'**

	<i>what is happening now</i>	<i>what could happen in the future</i>
Amount of evidence	low	low
Level of agreement or consensus	moderate	moderate
Level of confidence	low	low

Our knowledge of gross status changes is good for seals but moderate-poor for cetaceans, whilst our understanding of links between demography and climate for both groups is poor. We therefore can only have low level of confidence in what is happening now in relation to climate. As for what may happen in the future, without any clear understanding of impacts on marine mammals, in most cases it is impossible to make confident predictions.

Some marine mammal scientists are more confident than others that we are witnessing ecological effects of climate change as opposed to responses of individuals and local populations to local environmental variability. The statistical power to discriminate between the two remains low. Although more could be done to improve the current level of confidence, there will always be an upper limit to our ability to link population changes to the physical or biological drivers associated with climate change, particularly for some of the less accessible cetacean species.

**Can you comment on specific knowledge gaps which need addressing in this field**

- A pre-requisite to assessing impacts of climate change on marine mammals at a population level is a long-term, wide-ranging, monitoring programme that can discriminate between regional population responses and those occurring on a wider geographical scale. This is presently lacking for all UK cetacean species, whilst for seals there remain regions (e.g. Irish Sea) with only patchy coverage.
- Our knowledge of trends in basic life history parameters (growth rates, age at sexual maturity, reproductive rates, and mortality) for all cetacean species with the possible exception of harbour porpoise is woefully inadequate, based upon small sample sizes from a restricted number of areas, and without long-term continuity of data. For the majority of species, we are unlikely to obtain adequate information in the foreseeable future given how difficult they are to study and the resources available to do so. However, certain species could be targeted for more intensive study with some likelihood of success. In UK waters, these include (in addition to harbour porpoise) bottlenose dolphin, short-beaked common dolphin, white-beaked dolphin, Risso's dolphin, and minke whale.

- Functional responses to environmental change through physiological and behavioural mechanisms are also poorly understood for most marine mammal species. For this, seals are rather better suited to experimental studies where variables can be controlled. Once individual responses are better understood, it may be possible to make predictions at the population level.
- Too little is known about how changes in fish, cephalopod and plankton dispersion, distribution and abundance may affect the foraging ecology of particular marine mammal species. Often, one of the major gaps in information lies in the lack of data for non-commercial fish and cephalopod species, although even for some commercial species, such information is lacking..

**What are the major commercial effects that the impacts of climate change will have in this field?**

No marine mammal species in UK is exploited directly, although changes in the status and distribution of marine mammals could potentially have commercial effects if species that are targeted by the ecotourism industry become scarce, or alternatively if there are changes in competitive relations (e.g. an increase in seal predation upon commercially important fish).

**Please list full references (follow the conventions described below). Also provide links to other sources of information, websites and datasets that are of relevance to your submission.**

- Arnott, S. A. and Ruxton, G. D. (2002). Sandeel recruitment in the North Sea: demographic, climatic and trophic effects. *Marine Ecology Progress Series*, **238**, 199-210.
- Baines, M.E., Anderwald, P., and Evans, P.G.H. (2006) Monitoring a changing world - searching the past for long-term trends in the occurrence of cetaceans around the UK. *European Research on Cetaceans*, **20**: 1-11.
- Bakker, J. and Smeenk, C. (1987) Time-series analysis of *Tursiops truncatus*, *Delphinus delphis*, and *Lagenorhynchus albirostris* strandings on the Dutch coast. Pp. 14-19. In *The European Cetacean Society Report of the 1987 Meeting*, Hirtshals, Denmark, 26-28 January 1987 (Eds. J.W. Broekema & C. Smeenk).
- Beare, D., Burns, F., Greig, A., Jones E.G., Peach, K., Kienzle, M., McKenzie E., and Reid, D.G. (2004). Long-term increases in prevalence of North Sea fishes having southern biogeographic affinities. *Marine Ecology Progress Series*, **284**: 269-278.
- Beaugrand, G., Reid, P.C., and Ibanez, F. (2002). Reorganization of North Atlantic marine copepod biodiversity and climate. *Science*, **296**: 1692-1694.

- Boyle, P. (Ed.) (1983). *Cephalopod Life Cycles*. Academic Press, London.
- Brander, K., Blom, G., Borges, M.F., Erzini, K., Henderson, G., MacKenzie, B.R., Mendes, H., Ribeiro, J., Santos, A.M.P., and Toresen, R. (2003). Changes in fish distribution in the eastern North Atlantic: Are we seeing a coherent response to changing temperature? *ICES Marine Science Symposia*, **219**: 261-270.
- Bjørge, A. (2002). How persistent are marine mammal habitats in an ocean of variability? Pp. 63-91. In: Evans, P.G.H. and Raga, J.A. (Eds) *Marine mammals – biology and conservation*. Kluwer Academic/Plenum Publishers, New York.
- Camphuysen, K. and Peet, G. (2006). *Whales and Dolphins of the North Sea*. Fontaine Uitgevers, Graveland /The North Sea Foundation, Utrecht, The Netherlands. 159pp.
- Derocher, E., Lunn, N., and Stirling, I. (2004). Polar bears in a warming climate, *Integrative and Comparative Biology*, **44**; 163-176.
- Domingo, M., Kennedy, S., and Van Bresselem, M-F. (2002). Marine Mammal Mass Mortalities. Pp. 425-456. In: Evans, P.G.H. and Raga, J.A. (Eds) *Marine mammals – biology and conservation*. Kluwer Academic/Plenum Publishers, New York.
- Evans, P.G.H., Anderwald, P., and Baines, M.E. (2003). *Status Review of UK Cetaceans*. Report to English Nature & Countryside Council for Wales. 160pp. (obtainable from Sea Watch Foundation, 11 Jersey Road, Oxford OX4 4RT).
- Evans, P.G.H., Lockyer, C.H., and Read, A. J. (2007). Harbour Porpoise *Phocoena phocoena*. In: *New Handbook of British Mammals* (Editor D.W. Yalden). Mammal Society, London. (In press).
- Ferguson, S., Stirling, I., and McLoughlin, P. (2005). Climate change and ringed seal (*Phoca hispida*) recruitment in western Hudson Bay, *Marine Mammal Science*, **21**(1); 121-135.
- Forcada, J., Trathan, P., Reid, K., and Murray, E. (2005). The effects of global climate variability in pup production of Antarctic fur seals, *Ecology*, **86**(9); 2408-2417.
- Fraser, F.C. (1946). *Report on Cetacea stranded on the British Coasts from 1933 to 1937*. The British Museum (Natural History), London. 56pp.
- Frederiksen, M., Harris, M.P., Daunt, F., Rothery, P., and Wanless, S. (2004). Scale-dependent climate signals drive breeding phenology of three seabird species. *Global Change Biology*, **10**, 1214-1221.

- Frederiksen, M., Wright, P.J., Harris, M.P., Mavor, R.A., Heubeck, M., and Wanless, S. (2005). Regional patterns of kittiwake *Rissa tridactyla* breeding success are related to variability in sandeel recruitment. *Marine Ecology Progress Series*, **300**: 201-211.
- Geraci, J. and Lounsbury, V. (2002). Marine mammal health: holding the balance in an ever changing sea. Pp. 365-384. In: Evans, P.G.H. and Raga, J.A. (Eds) *Marine mammals – biology and conservation*. Kluwer Academic/Plenum Publishers, New York.
- Geraci, J., Harwood, J., and Lounsbury, V. (1999). Marine mammal die-offs. Pp. 367-395. In: Twiss Jr, J.R. & Reeves, R.R. (Eds.) *Conservation and Management of Marine Mammals*. Smithsonian Institution Press, Washington D.C.
- Greene, C.H. and Pershing, A.J. (2004). Climate and the conservation biology of North Atlantic right whales: the right whale at the wrong time? *Frontiers in Ecology and the Environment*, **2**(1): 29-34.
- Harvell, C.D., Mitchell, C.E., Ward, J.R., Altizer, S., Dobson, A.P., Ostfield, R.S., and Samuel, M.E. (2002). Climate warming and disease risks for terrestrial and marine quota. *Science*, **296**: 2159-2162.
- Harwood, J. (2001). Marine mammals and their environment in the twenty-first century. *Journal of Mammalogy*, **82**: 630-640.
- Heide-Jorgensen, M.P. and Laidre, K.L. (2004). Declining extent of open water refugia for top predators in Baffin Bay and adjacent waters, *Ambio*, **33**(8): 487-94.
- Hernández, M., Robinson, I., Aguilar, A., Gonzalez, L.M., López-Jurado, L.F., Reyero, M.I., Cacho, E., Franco, J., López-Rodas, V., and Costas, E. (1998). Did algal toxins cause monk seal mortality? *Nature*, **393**: 28.
- Hulme, M., Jenkins, G.J., Lu, X., Turnpenny, J.R., Mitchell, T.D., Jones, R.G., Lowe, J., Murphy, J.M., Hassell, D., Boorman, P., McDonald, R., and Hill, S. (2002). *Climate change scenarios for the United Kingdom: the UKCIP02 scientific report*. Tyndall Centre for Climate Change Research, Norwich.
- International Whaling Commission (1999). Report of the IWC workshop on climate change and cetaceans. *Report of the International Whaling Commission*, **47**: 293-313.
- Kinze, C.C., Addink, M., Smeenk, C., García Hartmann, M., Richards, H.W., Sonntag, R.P., and Benke, H. (1997) The white-beaked dolphin (*Lagenorhynchus albirostris*) and the white-sided dolphin (*Lagenorhynchus acutus*) in the North and Baltic Seas: review of available information. *Report of the International Whaling Commission*, **47**: 675-681.

- Lafferty, K.D., Porter, J.W., and Ford, S.E. (2004) Are diseases increasing in the ocean? *Annual Review of Ecology and Systematics*, **35**: 31-54.
- Laidre, K. and Heide-Jorgensen, M. (2005), Arctic sea ice trends and narwhal vulnerability, *Biological Conservation*, **121**: 509-517.
- Leaper, R., Cooke, J., Trathan, P., Reid, K., Rowantree, V., and Payne, R. (2005) Global climate drives southern right whale (*Eubaeana australis*) population dynamics, *Biology Letters*, **2**(2) (published online DOI:10.1098/rsbl.2005.0431).
- Learmonth J.A., Macleod C.D., Santos M.B., Pierce G.J., Crick, H.Q.P., and Robinson, R.A. (2006) Potential effects of climate change on marine mammals. *Oceanography and Marine Biology*, **44**: 431-464.
- MacDonald, R., Harner, T., and Fyfe, J. (2005). Recent climate change in the Arctic and its impact on contaminant pathways and interpretation of temporal trend data, *Science of the Total Environment*, **342**; 5-86.
- MacLeod, C., Bannon, S., Pierce, G., Schweder, C., Learmonth, J., Herman, J., and Reid, R. (2005). Climate change and the cetacean community of north-west Scotland. *Biological Conservation*, **124**: 477-483.
- MacLeod, C.D., Santos, M.B., Reid, R.J., Scott, B.E., and Pierce, G.J. (2007). Linking sandeel consumption and the likelihood of starvation in harbour porpoises in the Scottish North Sea: Could climate change mean more starving porpoises? *Biology Letters* (published online, DOI: 10.1098/rsbl.2006.0588).
- MacLeod, C.D., Weir, C.R., Pierpoint, C., and Harland, E.J. (2007). The habitat preferences of marine mammals west of Scotland (UK). *Journal of the Marine Biological Association of the United Kingdom*, **87**: 157-164.
- Mavor, R.A., Parsons, M., Heubeck, M., and Schmitt, S. (2006). *Seabird numbers and breeding success in Britain and Ireland, 2005*. Peterborough, Joint Nature Conservation Committee. (UK Nature Conservation, No.30).
- McMahon, C. and Burton, C. (2005). Climate change and seal survival: evidence for environmentally mediated changes in elephant seal *Mirounga leonina* pup survival, *Proceedings of the Royal Society B*, **272**; 923-928.
- Peperzak, L. (2003). Climate change and harmful algal blooms in the North Sea. *Acta Oecologica*, **24**, Supplement 1: 139-144.
- Perry, A.L., Low, P.J., Ellis J.R., and Reynolds J.D (2005). Climate Change and Distribution Shifts in Marine Fishes. *Science*, **308**: 1912-1915.

- Pörtner, H.O., Langenbuch, M., and Reipschläger, A. (2004). Biological impact of elevated ocean CO<sub>2</sub> concentrations: lessons from animal physiology and earth history. *Journal of Oceanography*, **60**: 705-718.
- Reid, C.R., Edwards, M., Beaugrand, G., Skogen, M., and Stevens, D. (2003). Periodic changes in the zooplankton of the North Sea during the twentieth century linked to oceanic inflow. *Fisheries Oceanography*, **12**: 260-269.
- Rindorf, A., Wanless, S., and Harris, M. P. (2000). Effects of changes in sandeel availability on the reproductive output of seabirds. *Marine Ecology Progress Series*, **202**: 241-252.
- Rose, G.A. (2005). On distributional responses of North Atlantic fishes to climate change. *ICES Journal of Marine Science*, **62**: 1360-1374.
- Royal Society (2005). *Ocean acidification due to increasing carbon dioxide*. Policy Document 12/05. The Royal Society, London. 60pp.
- Santos, M.B., Pierce, G.J., Learmonth, J.A., Reid, R.J., Ross, H.M., Patterson, I.A.P., Reid, D.G., and Beare, D. (2004) Variability in the diet of harbor porpoises (*Phocoena phocoena*) in Scottish waters 1992-2003. *Marine Mammal Science*, **20**: 1-27.
- Scholín, C.A., Gulland, F., Doucette, G.J., Benson, S., Busman, M., Chavez, F.P., Cordaro, J., DeLong, R., De Vogelaere, A., Harvey, J., Haulena, M., Lefebvre, K., Lipscomb, T., Loscutoff, S., Lowenstine, L.J., Maein III, R., Miller, P.E., McLellan, W.A., Moeller, P.D.R., Powell, C.L., Rowles, T., Silvagni, P., Silver, M., Spraker, T., Trainer, V., and Van Dolah, F.M. (2000). Mortality of sea lions along the central California coast linked to a toxic diatom bloom. *Nature*, **403**: 80-84.
- Sheldrick, M.C. (1989). Stranded whale records for the entire British coastline, 1967 – 1986. *Investigations on Cetacea*, **22**: 298-329.
- Short, F.T. and Neckles, H.A. (1999). The effects of global climate change on seagrasses. *Aquatic Botany*, **63**: 169-196.
- Simmonds, M. and Isaac, S. (2007). The impacts of climate change on marine mammals: early signs of significant problems. *Oryx*, **41**: 1-8.
- Stenseth, N.C., Mysterud, A., Otterlsen, G., Hurrell, J.W., Chan, K.-S., and Lima, M. (2002). Ecological effects of climate fluctuations. *Science*, **297**: 1292-1296.
- Stenseth, N. C., Ottersen, G., Hurrell, J. W., and Belgrano, A. (2004). *Marine ecosystems and climate variation*. Oxford University Press, Oxford.
- Stirling, I., Lunn, N.J., and Iacozza, J. (1999). Long-term trends in the population ecology of polar bears in western Hudson Bay in relation to climate change. *Arctic*, **52**: 294-306.

- Sun, L., Liu, X., Yin, X., Zhu, R., Xie, Z., and Wang, Y. (2004). A 1,500-year record of Antarctic seal populations in response to climate change *Polar Biology*, **27**: 495-501.
- Trites, A., Miller, A., Maschner, H., Alexander, M., Bograd, S., Calder, J., Capotondi, A., Coyle, K., Di Lorenzo, E., Finney, B., Gregr, E., Grosch, C., Hare, S., Hunt, G., Jahncke, J., Kachel, N., Kim, H., Ladd, C., Mantua, N., Marzban, C., Maslowski, W., Mendelssohn, R., Neilson, D., Okkonen, S., Overland, S., Reedy-Maschner, K., Royer, T., Schwing, F., Wang J., and Winship, A. (2006). Bottom up forcing and decline of Stellar Sea Lions in Alaska: assessing the ocean climate hypothesis, *Fisheries Oceanography*.
- Tynan, C. and DeMaster, D. (1997). Observations and predictions of arctic climatic change: potential effects on marine mammals. *Arctic*, **50**: 308-322.
- Walther, G., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T., Fromentin, J., Hoegh-Guldberg, O., and Bairlein, F. (2002). Ecological responses to recent climate change. *Nature*, **416**; 389-395.
- Wanless, S., Harris, M.P., Redman, P., and Speakman, J.R. (2005). Low energy values of fish as a probable cause of a major seabird breeding failure in the North Sea. *Marine Ecology Progress Series*, **294**: 1-8.
- Würsig, B., Reeves, R.R., and Ortega-Ortiz, J.G. (2002). Global climate change and marine mammals. Pp. 589-608. In: Evans, P.G.H. and Raga, J.A. (Eds) *Marine mammals – biology and conservation*. Kluwer Academic/Plenum Publishers, New York.